

Prevention - The most effective way of preventing measles is through the MMR immunisation. Most children in the UK receive the MMR immunisation.

Children born on or after 1st July 2024 will receive their **first dose at 12months** and their **second dose at 18 months**.

Children born before 1st July 2024 will receive their **first dose at 12months** and their **second dose at 3 years and 4 months**. One dose of a measles containing vaccine is about 90 per cent effective against measles. A second dose of MMR was introduced to the UK vaccination schedule in the mid 1990s, before children started school, which is very important as it raises their immunity against the disease to almost 100 per cent.

Herd immunity - Vaccination is important.

As measles is so infectious, a high level of immunity is needed before 'herd immunity' can be reached. Herd immunity occurs when a certain percentage of a population is vaccinated and the spread of the disease is effectively stopped. To reach this level of protection, about 85 percent of preschool children should be immune (for this to happen, 90-95 per cent need to be immunised with at least one dose of vaccine).

In 2024, only 85% of people in the UK had received two doses of the MMR vaccine. This is significantly lower than the 95% target recommended by the World Health Organisation (WHO) to achieve herd immunity and prevent outbreaks.

Whiston Hospital
Warrington Road,
Prescot, Merseyside, L35 5DR
Telephone: 0151 426 1600

St Helens Hospital
Marshalls Cross Road,
St Helens, Merseyside, WA9 3DA
Telephone: 01744 26633

Southport Hospital
Town Lane, Kew,
Southport, Merseyside,
PR8 6PN
Telephone: 01704 547 471

Ormskirk Hospital
Dicconson Way, Wigan Road,
Ormskirk, Lancashire, L39 2AZ
Telephone: 01695 577 111



**Mersey and West Lancashire
Teaching Hospitals**
NHS Trust

Measles

Patient information leaflet

If you need this leaflet in a different language or accessible format please speak to a member of staff who can arrange it for you.

اگر به این بروشور به زبان دیگر یا در قالب دسترس پذیر نیاز دارید، لطفاً با یکی از کارکنان صحبت کنید تا آن را برای شما تهیه کند.

Jeśli niniejsza ulotka ma być dostępna w innym języku lub formie, proszę skontaktować się z członkiem personelu, który ją dla Państwa przygotowuje.

Dacă aveți nevoie de această broșură într-o altă limbă sau într-un format accesibil, vă rog să discutați cu un membru al personalului să se ocupe de acest lucru pentru dumneavoastră

如果您需要本传单的其他语言版本或无障碍格式，请联系工作人员为您安排。

إذا احتجت إلى هذه النشرة بلغة أخرى، أو بتيسيق يسهل الوصول إليه، يرجى التحدث إلى أحد الموظفين لترتيب ذلك لك.

Author: Nurse
Department: Paediatrics
Document Number: MWL2777
Version: 007
Review Date: 30/09/2028

The nature of the condition - Few parents today will remember the potentially devastating effects of measles before the introduction of a vaccine against this disease.

In fact, complications were relatively common, and measles claimed the lives of around 85 children every year in the UK (more in epidemic years) before the vaccine became part of the routine immunisation schedule in 1968.

Fortunately the vast majority of children in the UK are now protected against measles. A huge amount of research worldwide has overwhelmingly confirmed the safety of the MMR vaccine, and has led to a return in public confidence following scare stories raised by one doctor in the late 90s. But in some areas where uptake of the vaccine is low, there have recently been outbreaks of measles.

In 2023, approximately 10.3 million people were infected with measles globally, primarily among children under five.

In England, for the period January to June 2023, 33 out of 91 (36%) reported measles cases were in children under five years old. 17.5% of hospitalised measles patients were admitted to the paediatric intensive care unit (PICU) for severe complications like respiratory failure. So it is worth knowing a bit about measles – what to look out for, and how the vaccine can protect your child.

Measles is a highly contagious viral infection. It spreads easily from an infected person via coughs and sneezes, and even just by breathing.

Symptoms develop between six and 19 days after contact with the infection. Children with measles are usually infectious themselves for one or two days before the rash appears – in other words, before it is confirmed that they have measles.

The first symptoms, which usually last two or three days, include feeling generally unwell and having a runny nose, cough, sore eyes, and a fever which may be as high as 40C. Children may also complain that bright light hurts their eyes.

A day or two before the rash appears, some children develop 'Koplik's spots' – tiny white spots on a red background inside the mouth. The rash appears on the fourth day of the illness and is red and blotchy.

It starts at the hairline and travels down the body over a period of about three days, although it is mainly confined to the face and upper body. After three to four days, the rash turns a brownish colour and gradually fades. Having measles is believed to give a child lifelong immunity against the infection.

The condition does have what is known as an 'immunosuppressive' effect – so for a period of months after the attack, your child might be more susceptible than usual to other infections.

Complications - Most children with measles make a full recovery – but unfortunately, around 1 in 15 suffer from complications. These include pneumonia, middle ear infections and convulsions.

Making a diagnosis - The measles rash is very distinctive, but it is important to see your GP. Your doctor will take a sample of your child's saliva for testing to confirm the diagnosis.

Measles is a notifiable disease – each case needs to be reported to the Health Protection Agency which your GP or hospital doctor will be able to do. It is best to call the surgery or hospital beforehand to warn them your child might have measles, as there may be a separate waiting area for people with infectious diseases.

Benefits of treatment - There is not any specific treatment for measles itself, only the symptoms. You are advised to ensure your child has plenty to drink, offer paracetamol to reduce fever and contact the GP surgery or hospital if you suspect your child's condition is deteriorating.

Measles can be very dangerous to the unborn child. It is really important to keep a child with measles away from anyone who is pregnant during the infectious period, as the woman may not have had measles.