

Medial Branch Block of Lumbar Facet Joint

If you need this leaflet in a different language or accessible format
please speak to a member of staff who can arrange it for you.

اگر به این بروشور به زبان دیگر یا در قالب دسترسی‌بذرگ‌دارید،
لطفاً با یک از کارکنان صحبت کنید تا آن را برای شما تهیه کند.

Jeśli niniejsza ulotka ma być dostępna w innym języku lub formacie,
proszę skontaktować się z członkiem personelu, który ją dla Państwa przygotuje.

Dacă aveți nevoie de această broșură într-o altă limbă sau într-un format accesibil,
vă rog să discutați cu un membru al personalului să se ocupe
de acest lucru pentru dumneavoastră

如果您需要本传单的其他语言版本或无障碍格式, 请联系工作人员为您安排。

إذا احتجت إلى هذه النشرة بلغة أخرى، أو بتنسيق
يسهل الوصول إليها، يرجى التحدث إلى أحد الموظفين لترتيب ذلك لك.

Southport & Formby Hospital
Town Lane, Kew,
Southport, Merseyside,
PR8 6PN
Telephone: 01704 547471

Ormskirk Hospital
Dicconson Way, Wigan Road,
Ormskirk, Lancashire, L39 2AZ
Telephone: 01695 577 111

www.MerseyWestLancs.co.uk

Author: Specialist Nurse
Department: Pain Management Southport & Ormskirk Hospital
Document Number: 388 (2)
Version: 1
Review Date: 30 / 11 / 2026

The aim of this procedure is to lessen the experience of pain. Time in the pain management clinic is available for you to discuss possible options available with the doctor providing this treatment. Prior to treatment you will be asked to sign a consent form.


Potential risks are listed on the consent form for your information so that you may read and understand the treatment you are about to receive.

What is a Facet Joint Medial Branch Block?

The facet joints are the small linking joints of the spine. At each level of the spine, nerve supply to the facet joints is provided by small nerves emerging from the spinal cord. These are called the medial branch nerves. They supply sensation from the facet joint.

Many people experience pain in these joints, which may be linked to arthritic changes or general 'wear and tear' which we all experience as we get older.

The picture explains where the facet joints are in the spine.



Special Instructions

Please do not drive for the next 24 hours after receiving this injection.

Any Condition Specific Danger Signals to Look Out For: See risk/side effects section

Contact Information If You Are Worried About Your Condition

GP

Other Useful Telephone Numbers / Contacts:

NHS 111
Stop Smoking Helpline (Sefton) - 0300 100 1000
Stop Smoking Helpline (West Lancashire) - 0800 328 6297

Contact Details

Chronic Pain Management Contact Details

(01704) 704201

Monday - Wednesday

Chronic Pain Management Contact Details

(01695) 656421

Wednesday – Fridays

Appointments office

01695 656680

Daycase Appointments

01695 656007

Specialist Nurse and Pain Nurses

Telephone: 01704 705132



**Please use QR Code to
complete the Patient
Satisfaction Survey**

This procedure is carried out in an appropriate treatment area either under ultrasound or X-ray screening to aid accuracy. A solution of local anaesthetic medication and a slow release steroid preparation is injected around the nerve supplying sensation to the facet joint.

The skin is made numb with local anaesthetic before the medial branch block is performed. The steroid preparation has an anti-inflammatory action and will remain in your system for up to 12 weeks.

This does not denote the time the effects of the injection last for: you may experience longer or shorter pain relief than this, as every person is different. Many people experience significantly longer pain relief.

To receive the injection you will be required to lie face down on the table throughout the procedure. There may be some discomfort at the time of the injection but this will not last long.

It is safe to take your normal medication and eat breakfast prior to attending the hospital.

Following the procedure you will be offered a hot drink. After consuming this, if you feel well enough you can travel home.

Please read your appointment card carefully as this will tell you the clinic area and the hospital site you must attend for this procedure.

Please ensure you do not drive for 24hrs after the procedure.

Risks/ Side Effects

All steroid injections can cause a short term reduction in the function of the immune system along with possible facial flushing, short term menstrual irregularity in females and altered blood sugar levels in diabetic patients. Further information on the risk can be found in the consent form.

Infection

There is a small chance of wound infection or the potential for deeper spinal infection. We do take every measure to avoid this and procedures are conducted in a theatre environment aseptically to reduce risk.

Bleeding

Rarely may you get some bleeding and or bruising around the injection site this will usually settle after a few days.

Nerve Injury

There is the rare potential for a nerve injury. This can happen if the needle comes into contact with a nerve or nerve sheath. This may cause some numbness and weakness affecting the limbs. This is generally temporary whilst the local anesthetic wears off.

Injection Site Tenderness

You may experience some injection site tenderness and increased pain for a few days or hours after injection. This generally settles on its own.

Alternative Treatment Options

If you decide not to have the procedure your alternative option is to continue to manage your pain conservatively.

This can be assisted with physiotherapy and referral to a pain management programme if suitable.

In some cases if appropriate and clinically indicated, you may choose to proceed with surgery instead.

The aims benefits and alternatives to a Medial Branch Injection have been outlined within the leaflet. If you decide you do not wish to go ahead with your planned procedure, please contact the Waiting List office on 01695 656670 to inform them of your decision as soon as possible.

On the day of your injection, you and your doctor will sign the consent form together to document your consent to proceed with the procedure.

This patient information leaflet is intended to be used to support discussion during your clinical consultation. If there is anything you do not understand or are unsure about, please ask the doctor at your appointment or contact the people below.

You will be given a follow-up appointment before you leave the clinic